


Sortni poskus/Variety trial
Maribor

Prejšnji posevek: oljne buče
Gnojenje: 122 kg/ha N (v 4 obrokih), 72 kg/ha P₂O₅, 140 kg/ha K₂O
Škropljenje: Boxer 2,5 l/ha, Bizon 2,5 l/ha, Ascra Xpro 1,5 l/ha, Elatus era 1 l/ha
Gostota setve: medvrstna razdalja 12,5 cm, 400-750 kalivih zrn/m²
Setev: 24.10.2021, vznik: 9.11.2021, žetev: 12.07.2022
Zasnova poskusa: naključni blok v 4 ponovitvah, osnovna parcela 7,5 m²

Ozimna rž/Winter rye

Previous crop: oil pumpkins
Fertilization: 122 kg/ha N (in 4 rations), 72 kg /ha P₂O₅, 140 kg /ha K₂O,
Spraying: Boxer 2,5 l/ha, Bizon 2,5 l/ha, Ascra Xpro 1,5 l/ha, Elatus era 1 l/ha
Sowing density: interrow spacing 12.5 cm, 400-750 germinative seeds/m²
Sowing: 24.10.2021, emergence: 9.11.2021, harvesting: 12.07.2022
Trial layout: randomized block design in 4 repetitions, plot size 7,5 m²

 SORTA VARIETY	Datum klasitve <i>Date of earing</i>	RASTLINA/PLANT		ANALIZA/ANALYSIS				PRIDELEK/YIELD			BOLEZEN/DISIASE				
		Višina <i>Height</i>	Poleg po klasitvi <i>Lodging from ear forming on</i>	Hektolitrska masa <i>Test weight</i>	Absolutna masa zrnja <i>Thousand seeds weight</i>	Surove beljakovine <i>Crude protein</i>	Število padanja <i>Falling number</i>	Vlaga zrnja <i>Grain moisture</i>	Zrnje s 14% vlago <i>Grain with 14% moisture</i>	Rodnostna skupina <i>Yield category</i>	<i>Erysiphe graminis</i>	<i>Puccinia graminis</i>	<i>Puccinia recondita</i>	<i>Rhynchosporium secalis</i>	<i>Fusarium sp.</i>
		cm	1-9 1=brez/without	kg	g	%	s	%	t/ha		1-9 1=brez/without	1-9 1=brez/without	1-9 1=brez/without	1-9 1=brez/without	1-9 1=brez/without
KWS JEHTRO (H)	0. 1.	147	4	74	46	8,8	327	15,1	13,12	I	1	1	1	3	1
SU FORSETTI (H)	0. 1.	139	5	74	45	10,1	301	15,5	12,63	II/1	1	1	1	3	1
ASTRANOS (H)	0. 1.	142	4	74	47	11,6	296	11,5	12,51	II/1	1	1	1	3	1
HELLTOP (H)	0. 1.	150	5	73	43	9,7	264	12,0	11,30	II/3	1	1	1	4	1
DANKOWSKIE AMBER (POP)	0. 1.	140	7	74	41	9,2	296	12,4	9,99	III	1	1	1	3	1
Povprečje/Mean									11,91						
LSD (0,05)									1,14						